



Attractive contrast of shiny flowers above dark leaves



1. Home

Begonia semperflorens F?

Nightlife

• Traditional compact, bronze leaved semperflorens



- Early flowering
- Perfectly works in pots and packs
- Dark foliage partner to Sprint Plus
- Attractive contrast of shiny flowers above dark leaves

Bookmark

Recommend

Print

Crop Time

Spring: 10 - 12 weeks

Height

9?/23 cm

Exposure Sun - Shade

Seed Form

Raw Seed, Pelleted Seed

Best Uses Bedding, Landscape

Culture guide

Usage

Annual, packs, pots, containers and landscape plantings

Sow time

December onwards

Sowing method



1-2 seeds/pills per plug

Germination

Light is required for germination. 5-10 days at temperatures of 75-80 °F (24-26 °C), and relative humidity levels of 95% or higher. For pelleted seed, slightly higher temperatures and 100% relative humidity aid in uniformity of germination. pH levels of 5.5-6.5 are optimum.

Growing on

Transplant plugs after 7-9 weeks and grow on at 62-68 °F (16-20 °C). Fertilize weekly with 150 ppm nitrogen watching to avoid excessive salt buildups.

Media

Use a well-drained, growing substrate, pH: 5.5-6.2.

Temperature

Grow at 17-19 °C. For selling harden the plants at 7-10 °C and UV light. Begonia semperflorens does not tolerate frost.

Fertilization

Moderate fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen, using a complete balanced fertilizer. High ammonium and high nitrogen levels cause big foliage. Avoid pH above 6.5, as high pH causes iron deficiency. Apply Fe-chelate if chlorosis becomes a problem. Avoid high salt levels in substrate.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.



Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

Download